

Exoplanets

Lecture 11
MFF UK
15.12.2020

Exoatmospheres

Outline

- Recap exoatmospheres
- Detection methods
 - spectroscopic
 - spectrophotometric
- Weather on exoplanets
- Challenges of precise spectroscopy/photometry
(J. Subjak input)

What do we know?

- Atmospheres of exoplanets do exist
- We know different types of atmospheres on exoplanets
 - H/He rich, heavy elements rich, water
 - the thinner the atmosphere is the more challenging is its detection
- Large telescopes with precise instruments needed, but....

Scale height

$$H = \frac{kT}{Mg}$$

k – Boltzmann constant

M – mean molecular weight





g – gravitational constant

T – mean atmospheric temperature

EARTH – about 8 km

TITAN - about 40 km


Different types of atmospheres



And how to detect the atmospheres?

- After new detections of exoplanets, also characterization attempts start in 2002
- Main goals are:
 - detection of atmosphere
 - physical conditions on the surface/in the atmosphere of the exoplanet
- Photometric and spectroscopic methods

Transits and eclipses of exoplanets




From Angerhausen et al. 2008

Transit spectroscopy, the principle

Transit spectroscopy = transmission spectroscopy


$$\text{Signal} = \text{Annulus} / R_{\text{star}}^2$$



Typical Signal of the planetary spectral lines $< 10^{-4}$


Smaller star & larger planet = better chance to see something

Transmission spectroscopy high spectral resolution



HD209458 b

- Charbonneau et al. 2002
<https://arxiv.org/pdf/astro-ph/0111544.pdf>
- Detection with HST STIS



What can we see?

- Absorption in stellar lines due to planetary atmosphere by atoms – high. resolution spectroscopy (Na, K)
- Absorption in stellar lines due to planetary atmosphere by molecules – low. resolution spectroscopy (H₂O, CO₂, TiO, CH₄)
- First observations performed in 2002 with HST
 - HD209458b





Figure 1. **Left:** The Lyman α stellar line as observed by Vidal-Madjar et al. (2003). The averaged profile observed during transit (thick line) presents a reduced flux when compared to the pre-transit profile (thin line). The region named “Geo” corresponds to the region where the geocoronal Lyman α correction was too important. In the “In” region absorption is observed while the “Out” region serves as a flux reference. **Right:** The averaged “In”/“Out” flux ratio in the individual exposures of the three observed transits (see text). Exposures A1, B1, and C1 were performed before and A2, B3, and C3 entirely during transits. Error bars are $\pm 1\sigma$. The “In”/“Out” ratio decreases by $\sim 15\%$ during the transit. The thick line represents the absorption ratio modeled through a particle simulation (see Fig. 3).

First ground based detection

- Redfield et al. 2008 - <https://iopscience.iop.org/article/10.1086/527475/pdf>
- Sodium doublet in HD189733b
- HET – 9.2-m telescope









Image from Redfield et al 2008

First ground-based detection with 4m class telescope


- Wyttenbach et al 2015 -
<https://arxiv.org/abs/1503.05581>
- HD187933b with HARPS



Another Sodium Detection with HARPS



Another Sodium Detection with HARPS



Another Sodium Detection with HARPS




Figure 4. The radial velocities of our targets have been measured with HARPS: WASP-76 (top left), WASP-166 (top right), WASP-127 (bottom left) and KELT-11 (bottom right). There is some apparent offset between various epochs for the same target, which may be due to some instrumental shift or could be caused by the planet crossing star spots or faculae regions. In each case, the orbital motion is also indicated with the dashed line. Any deviation from this is due to the Rossiter-McLaughlin effect, which is clearly in all but one of our targets.

Wasp-166b revisited




Fig. 4. HARPS sodium doublet transmission spectrum of WASP-166b for all nights combined shown in the PRF. Upper panel: In grey, data points at full HARPS resolution. In black, grey data points binned by $\times 25$ for visibility. The theoretical line centres for the sodium doublets are shown as vertical blue dashed lines, a Gaussian fit to the unbinned data is shown in red. The sodium absorption is visible for both lines of the doublet. The combined line contrast is measured at $0.455 \pm 0.135 \%$, resulting in a 3.4σ detection, see Sec. 4.3. Lower panel: Residuals of the Gaussian fit in %.

Wasp-166 revisited




Fig. A.1. The upper panel shows all spectra in the SRF as a 2D map of wavelength and transit phase for the first transit. The stellar sodium doublet is visible as two horizontal light yellow bands. Transit ingress and egress are marked with black dashed lines. The central plot shows the normalised sum of all spectra with a fit to each line in dashed blue. The FWHM is indicated as dotted vertical lines. The lower panel shows the same data, but corrected for the stellar spectrum by the master-out, in the PRF. The dotted lines propagate the position of the FWHM from the central panel. The low-SNR remnants are clearly visible, but the SNR is too low to see the planetary trace.

Wasp-127 revisited (with ESPRESSO)

R. Allart, L. Pino, C. Lovis et al.: WASP-127b seen by ESPRESSO




Fig. 8: Transmission spectrum around the Na D2 (*left*) and Na D1 (*right*) line averaged across the two transits in grey and binned by fifteen elements in black. The vertical blue dash dotted line represents the expected position of planetary sodium lines.

But... small telescopes?

- Kabath et al. 2019
- FEROS at 2-m telescope real data and injected sodium – DETECTION POSSIBLE!






Figure 4. Sanity test with injected signal: (left) the signature of sodium absorption in the planetary atmosphere was injected into the FEROS WASP-18b data set. The strength of the sodium signal was set to be equal to that detected from HD189733b by Wyttenbach et al. (2015). (right) Injected sodium signal of the equivalent strength to the WASP-49b detection by Wyttenbach et al. (2017). Black points in both panels correspond to binning by a factor 10. The red dashed lines indicate the position of the NaD lines.

Prospects for small telescopes




Figure 5. The expected atmospheric signal for well-characterised transiting planet as a function of the visual magnitude of the host star. The orange triangle shows the position of HD 189733, while the red triangle is the rescaled value, assuming a 2m-class telescope (see text). All the points above the solid line are suitable candidates to perform transmission spectroscopy with a 2m-class telescope.





Figure 6. Expected signal as in previous Fig. 5 but from TESS planets which are depicted as black dots. The orange line is the detection limit for 2-m telescopes and good candidates for further follow-up will be above the line (see text for more detail).

What happens if the star is flaring?


- Klocova 2017 et al. - <https://arxiv.org/pdf/1707.09831.pdf>
- What happens if during the transit a flare occurs




Other methods

Spectrophotometry

- Spectroscopy during the transit/eclipse
- Usually, low spectral resolution
- Spectral bins are selected to obtain spectrophotometric light curve (by integrating of the flux)
- Resulting light curve is fitted and transit parameters are obtained
- Depth of transit varies with wavelength
= TRANSMISSION SPECTRUM



Atmosphere spectrum




Exoplanet Transit Event



Towards Earth

FORS2 2010, 2011





Bean et al. 2010, Nature

Bean, Desert, Kabath et al. 2011,


SOFI NIR transmission spectroscopy


1.5 – 2.3 micron low res.
3 nights in 2011



WASP-19b – better resolution

- Sedaghati et al. 2015, A&A










Fig. 2. Transmission spectrum of WASP-19b as measured with FORS2 (black dots, with error bars) compared to two models of planetary atmospheres, one with no TiO (top panel) and one with a solar abundance of TiO (bottom panel), from Burrows et al. (2010) and Howe & Burrows (2012). We have also estimated the mean value of the models in bin sizes of 20 nm (orange open squares). The dashed lines represent the weighted mean and plus or minus three scale heights.

Can we determine the colour of skies on exoplanets?

- Rayleigh scattering - GJ3470b?






Very accurate photometry

Our observations with 4m class

- SOFI @ NTT – La Silla 3 nights
- OSIRIS @ SOAR - Cerro Pachon 1 night
- SOI @ SOAR - Cerro Pachon 1 night



Both telescopes are 4-m class!!!

ESO





SOAR

Our measurements - OSIRIS



$$R_p/R_s = 0.118101 (-)0.002766 (+)0.002562$$

Our measurements - SOI




SOAR I-BESSEL:

$$R_p/R_s = 0.117151 (-)0.001173$$
$$(+0.001182$$

Observations performed by S. Hoyer

4-m class telescopes good?



Our results compared (photometry)





Fig. 11. *Left:* A zoom-in from Fig. 10 for the optical region around our *I*-Bessel measurements. *Right:* The *K*-band region of spectra around our 2.14 μm observation. Our measurement points are represented by dark circles, while gray points follow the description in Fig. 10. A color version of this plot can be found in the electronic version of the paper.

Transits and eclipses of exoplanets



From Angerhausen et al. 2008

Emission from the planet

- Thermal radiation from the planet in IR



$$\text{Signal} = T_{\text{planet}}/T_{\text{star}}(R_{\text{planet}}/R_{\text{star}})^2$$

- Very shallow signals – few mmags
- Measuring directly the (missing) emission of the reflected light from the planet
- Result is an emission spectrum
- Due to geometry, not all planets hide behind the host star

Secondary eclipse photometry HD209458b

Měření: Spitzer 24 μ m

T_{pl} : ca 1130K





Deming et al. 2005 Nature

Secondary eclipse photometry from the ground

- Thermal radiation from the planet in IR

$$\text{Signal} = T_{\text{planet}}/T_{\text{star}} (R_{\text{planet}}/R_{\text{star}})^2$$


Typically few mmags for hot Jupiters



Gillon et al. 2009

Wavelength (μm)


TiO species absorbing the stellar heat?



Mandell et al. (2015), "Spectroscopic Evidence for a Temperature Inversion in the Dayside Atmosphere of the Hot Jupiter WASP-33b", arXiv:1505.01490

Weather on exoplanets


Eclipses/transits




From Winn, 2010, <http://arxiv.org/pdf/1001.2010v5.pdf>



Variation due to day/night cycle

- Near to mid IR with SPITZER (now no more possible)





Raining iron?



TOI-700 example models

ATMOSPHERIC MODELING OF TOI-700 D

7

#	Archetype	Surface type	Surface pressure (bar)	Primary atmospheric constituent	Minor constituents, partial pressures (bar)	Rotation	Surface temperature (K)	TOA albedo	Sea ice fraction (%)	Stratospheric water vapor (kg/kg)
1	Plain	aqua	1	N ₂	H ₂ O	synchronous	236.7	0.3968	76.2%	6.5014e-8
2	Modern Earth	aqua	1	N ₂	H ₂ O CO ₂ (4e-4) CH ₄ (1.7e-6)	synchronous	246.7	0.3893	71.2%	3.605e-7
3	Modern Earth	land	1	N ₂	H ₂ O CO ₂ (4e-4) CH ₄ (1.7e-6)	synchronous	232.7	0.2840	N/A	N/A
4	Archean Earth	aqua	1	N ₂	H ₂ O CO ₂ (0.01) CH ₄ (1.0e-4)	synchronous	258.4	0.3594	66.3%	7.316e-7
5	Archean Earth	aqua	1	N ₂	H ₂ O CO ₂ (0.1) CH ₄ (1.0e-3)	synchronous	263.2	0.3988	62.3%	2.644e-5
6	Archean Earth	aqua	1	N ₂	H ₂ O CO ₂ (0.1) CH ₄ (2.0e-3)	synchronous	263.7	0.3998	61.6%	3.645e-5
7	Archean Earth	land	1	N ₂	H ₂ O CO ₂ (0.1) CH ₄ (1.0e-3)	synchronous	251.5	0.2455	N/A	N/A
8	Archean Earth	aqua	0.5	N ₂	H ₂ O CO ₂ (0.1) CH ₄ (1.0e-3)	synchronous	256.8	0.3906	67.3%	5.615e-5
9	Archean Earth	aqua	4	N ₂	H ₂ O CO ₂ (0.1) CH ₄ (1.0e-3)	synchronous	307.2	0.317	0	6.406e-6
10	Archean Earth	aqua	10	N ₂	H ₂ O CO ₂ (0.1) CH ₄ (1.0e-3)	synchronous	360.8	0.2452	0	2.203e-6
11	Archean Earth	aqua	1	N ₂	H ₂ O CO ₂ (0.1) CH ₄ (1.0e-3)	2:1 resonant	325.7	0.131	0	6.095e-6
12	Archean Earth	land	1	N ₂	H ₂ O CO ₂ (0.1) CH ₄ (1.0e-3)	2:1 resonant	257.0	0.2547	N/A	N/A
13	H ₂ -supporting	aqua	1	N ₂	H ₂ O CO ₂ (0.1) CH ₄ (1.0e-3) H ₂ (0.1)	synchronous	267.6	0.394	60.3%	2.669e-5
14	Early Mars	aqua	0.5	CO ₂	H ₂ O	synchronous	266.2	0.3572	62.6%	2.802e-5
15	Early Mars	aqua	1	CO ₂	H ₂ O	synchronous	284.4	0.3775	0.05%	2.520e-5
16	Early Mars	aqua	2	CO ₂	H ₂ O	synchronous	324.3	0.2771	0%	3.027e-5
17	Early Mars	aqua	4	CO ₂	H ₂ O	synchronous	364.2	0.2214	0%	2.857e-4
18	Early Mars	land	1	CO ₂	none	synchronous	258.9	0.2304	N/A	N/A
19	Early Mars	land	4	CO ₂	none	synchronous	302.3	0.2165	N/A	N/A
20	Early Mars	land	10	CO ₂	none	synchronous	353.5	0.2310	N/A	N/A

Table 3. Global mean climatological properties for each of our simulations. By definition, sea ice fraction and stratospheric water vapor or not applicable for desiccated land planet cases. The stratospheric water vapor is taken at the model top, 0.01 mbar. All quantities given are

TOI-700 example models

ATMOSPHERIC MODELING OF TOI-700 D

11




Figure 4. Comparison of transmission spectra for different atmospheric archetypes. All simulations shown here are ocean-covered and have a surface pressure of 1 bar. Prominent features are labeled. The spectrum for the “hydrogen-supporting” atmosphere (10% H₂) has both the highest continuum and the largest 15 μm CO₂ feature.

TOI-700 example phase curves

18

SUISSA ET AL.






Figure 10. Phase curves for the majority of our simulations in this work. Aquaplanets are shown in the top panel; desiccated planets are shown in the bottom panel. Transit occurs at 180° . In the left panel, the contribution of reflected light dominates at wavelengths $< \sim 4.3 \mu\text{m}$, while thermally emitted light dominates at wavelengths $> \sim 4.3 \mu\text{m}$. In the right panel, the integrated fluxes for the phase curves are $15\text{--}21 \mu\text{m}$.


The Earthshine

- Sterzik et al. 2012, Nature
- Observing of the Earthsine reflected by the Moon
- FORS at VLT
- Polarization can help to describe the surface features


Observations	Observing date	
	25 April 2011, 09:00 ut	10 June 2011, 01:00 ut
View of Earth as seen from the Moon		
Sun–Earth–Moon phase (degrees)	87	102
Ocean fraction in Earthshine (%)	18	46
Vegetation fraction in Earthshine (%)	7	3
Tundra, shrub, ice and desert fraction in Earthshine (%)	3	1
Total cloud fraction in Earthshine (%)	72	50
Cloud fraction $\tau > 6$ (%)	42	27

The Earthshine


- The Atlantic side hints at decreasing PVI
- The Pacific side shows no correlation with PVI
- PVI is a Polarization Vegetation Index
- PVI reflects the difference of polarization between vegetated and not vegetated surfaces



Next week (Xmass session)
Life in the Universe
Habitable zones
Discussions (papers)



Kaltenegger, L. and Traub, W. (2009) Transits of Earth-Like Planets. *Astrophysical Journal*



JWST
Launch 2018
Ideal for characterization of small
planets in infrared
Image NASA